


Arkansas Audubon Society

2018 Field Reference to Arkansas Birds

Date _____ Time _____ Total Species Observed _____
 Location _____ County _____
 Observers _____



This publication provides information on 419 species of birds currently documented for Arkansas. It is based primarily on bird records maintained by the Arkansas Audubon Society, and secondarily on other sources, including *Arkansas Birds- Their Distribution and Abundance* (James and Neal, UA Press, 1986), Breeding Bird Surveys, Christmas Bird Counts, eBird, and published literature on Arkansas birds. Since the information that follows is generalized due to space constraints, it may not seem appropriate for a given species in some locales. Taxonomy and nomenclature follow the *Checklist of North American Birds*, Seventh Edition, AOU, 1998, as amended through the Fifty-ninth supplement, July 2018.

Approximate dates of arrival and departure are indicated by numbers 1-12 for the months of the year and letters A, B, C, D for the weeks of the month. A plus (+) sign immediately following indicates that discontinuous records outside the period are on file. For species that have nested in Arkansas, one of two symbols follows the name: an asterisk (*) means the species currently nests with regularity, either yearly, or at least, with some consistency; an open circle (°) means that the species does not nest regularly and that there are few or no such reports since 1950. Beneath the dates of arrival and departure are codes which categorize seasonal status, relative abundance, and, in some cases, the sections of the state of primary occurrence.

The months of the year are grouped into the following seasons:

- Sp-spring (March-May) S-summer (June-July)
 F-fall (August-November) W-winter (December-February)

In combination with these designations, the following abbreviations are used to denote seasonal status:

- P-Present, species seen in the state each month but not necessarily in all sections (Great Egret)
 M-Migrant, species passes through the state in spring and/or fall (Cape May Warbler)
 R-Resident, species remains for an extended period of time (Painted Bunting)
 L-Local, species occurs in only a few places (Great-tailed Grackle)
 V-Visitor, species usually not present for an extended period of time (Black-legged Kittiwake)
 Sa-Storm-assisted, species appears in the state in association with strong coastal storms; not usually present for an extended time (Sooty Tern)

The categories of relative abundance assume reasonable effort spent in suitable habitat during appropriate seasons and are defined as follows:

- c-common, always or almost always observed (Killdeer)
 fc-fairly common, seen on a majority of trips but usually in smaller numbers (Red-shouldered Hawk)
 u-uncommon, may be difficult to find; seen only on a minority of trips (Golden-winged Warbler)
 r-rare, may be seen annually or nearly so but not in all locales (Golden Eagle)
 vr-very rare, encountered at irregular or infrequent intervals (Ruff)
 occ-occasional, 3-4 state records overall with at least 2 records since 1950 (Long-tailed Jaeger)
 acc-accidental, 1-2 state records (Fork-tailed Flycatcher)
 irr-irruptive, variable in occurrence, number, and distribution (Red-breasted Nuthatch)

The abbreviations in parentheses following the statements of relative abundance indicate the primary but not necessarily the exclusive section of the state where the species occurs: (n) north, (s) south, (e) east, (w) west, (ce) central, (nc) north-central, (sc) south-central, (ec) east-central, (wc) west-central (ne) northeast, (se) southeast, (nw) northwest, and (sw) southwest.

| | | | |
|---|---|--|--------------------------|
| Order Anseriformes | | Duck, Tufted | WVacc (3, 2007) |
| Whistling-Ducks, Geese, Swans, and Ducks | | | |
| Whistling-Duck, Black-bellied* | 2D-11D+ SRu SRfc (sw, se, ce, wc) WVvr | Scaup, Greater | 10A-4B Mu WRu |
| Fulvous* | 4A-10C+ VLR SRLr (se) | Lesser | 10D-4D+ Mc WRc |
| Goose, Snow | 10A-4C+ Mc WRc (e,c) | Scoter, Surf | 10B-5B Mr WRvr |
| Ross's | 10D-4C+ Mfc WRc | White-winged | 10D-5D+ Mr Wvr |
| Greater White-fronted | 9D-4B+ Mfc WRfc (Lc e) | Black | 10D-3C Mvr WRvr |
| Bean-Goose, Tundra | WVacc (1, 2018) | Duck, Long-tailed | 11A-4A+ Mr Wvr |
| Brant | WVacc (1, 1990) | Bufflehead | 11A-4C+ WRc |
| Goose, Barnacle | WVacc (1, 2010) | Goldeneye, Common | 11B-4C WRLfc |
| Cackling | 11A-3D+ WRu (Lfc) | Barrow's | WVvr (6, 2014) |
| Canada* | P c | Merganser, Hooded* | P SRLu WRfc |
| Swan, Mute | 11D-2D+ WVvr | Common | 10A-3C+ WVr (n) |
| Trumpeter | 11A-5A WRLc SRLr | Red-breasted | 10D-4D+ Mu Wvr |
| Tundra | 11C-3D Mr WVvr (Lr nc) | Duck, Ruddy° | 10A-4C+ Mc WRc SRr |
| Duck, Wood* | P c | Order Galliformes Turkey and Quail | |
| Garganey | SpVacc (1, 1984) | Bobwhite, Northern* | P Lu |
| Teal, Blue-winged* | P Mc SRr (Lu nw) WRr (s) | Grouse, Ruffed | extirpated |
| Cinnamon | 2C-5B 9A-12B SpMr FMvr | Prairie-Chicken, Greater | extirpated |
| Shoveler, Northern | 8D-4C+ Mc WRc | Turkey, Wild* | P u |
| Gadwall | 9A-5D+ WRc | Order Podicipediformes Grebes | |
| Wigeon, Eurasian | FWVacc (2, 2013) | Grebe, Least | FVacc (1, 2008) |
| American | 9C-4C+ Mc WRc | Pied-billed* | P WRc SRLu |
| Mallard* | P WRc SRu | Horned | 10A-4A+ WRfc |
| Duck, American Black | 10A-3B+ WRLr (e) | Red-necked | 11D-4B WVvr |
| Mottled* | P SRLu (se, sw) WRLu (se) | Eared | 9A-5A+ Mu WRr (Lu se) |
| Pintail, Northern | 10A-4B+ Mc WRfc (e, sw) | Western | 10B-3D+ Mr Wvr |
| Teal, Green-winged | 9D-4C+ Mc WRc | Order Columbiformes Pigeons and Doves | |
| Canvasback | 10D-4D+ Mfc WRu | Pigeon, Rock* | P c |
| Redhead | 10D-4A+ Mfc WRu | Collared-Dove, Eurasian* | P c |
| Duck, Ring-necked | 10C-5B+ Mc WRc | Pigeon, Passenger | extinct |

| | | | | | | | |
|--|------------------------------|---|---------------------------------------|---------------------------------|---------------------------------|--|----------------------------|
| Dove, Inca* | P u Lfc (s, wc) | Rail, Virginia | 4A-5D 8D-2D+ MLu WRLr | Sanderling | 4C-5D 7C-10D+ SpMr FMfc | Gull, Little | 11D-3C+ WVvr |
| Ground-Dove, Common | 8C-5A Vr | Sora | 4A-5C 9A-12D+ Mfc WRr (cc, s) | Dunlin | 3B-5D 10A-12D+ Mfc WRr | Laughing | 7A-5B+ Vr |
| Dove, White-winged* | P Vr Lu (w) | Gallinule, Purple* | 4B-10C SRLr (cc, s) | Sandpiper, Purple | FVacc (1, 1976) | Franklin's | 3B-6A 9C-12C+ SpMu FMfc |
| Mourning* | P c | Common* | 4B-12D+ Mr (nw) SRLu (s) WRLr (sw) | Baird's | 3B-6A 7D-11C SpMu FMfc | Mew | WVacc (1, 2018) |
| Order Cuculiformes Cuckoos, Roadrunners, and Anis | | Coot, American* | P Mc WRc SRLr | Least | 7B-6B+ Mc WRu | Ring-billed | 8C-5A+ WRc |
| Cuckoo, Yellow-billed* | 4A-10C+ SRc | Crane, Sandhill | 10A-5A+ Mr WVr WRLu (sw, ne, e) | White-rumped | 4B-6C 8B-10D SpMfc FMvr | California | 12A-3B+ WVvr (wc) |
| Black-billed ^o | 4C-6D 8C-10C+ Mr SRvr (p) | Whooping | extirpated | Buff-breasted | 4A-5D 7D-10A+ SpMvr FMu | Herring | 8B-5C+ WRLu |
| Roadrunner, Greater* | P Lu (w, ce) Lr (e) | Order Charadriiformes Plover, Sandpipers, Gulls, and Terns | | Pectoral | 3A-6B 6D-11D+ Mc | Iceland | 11D-3A WVvr (wc) |
| Ani, Groove-billed | 5C-5D 9C-11D+ SpVacc FVvr | Stilt, Black-necked* | 3D-10D+ SRfc (cc, se) | Semipalmated | 4A-6B 7B-11A+ Mc | Lesser Black-backed | 10B-3B WVvr WRLr (wc) |
| Order Caprimulgiformes Goatsuckers | | Avocet, American | 4A-11D+ SpMu FMfc | Western | 4B-6A 7A-12D+ SpMu FMfc WRvr | Glaucous | 11D-3B WVvr |
| Nighthawk, Common* | 4B-11A+ SRc | Plover, Black-bellied | 3C-5D 8B-11D+ SpMu FMu | Dowitcher, Short-billed | 4D-6A 7D-10A Mu | Great Black-backed | WVacc (2, 2017) |
| Chuck-will's-widow* | 3D-9C SRc | Golden-Plover, American | 3A-5D 8C-11B+ SpMc FMu | Long-billed | 3A-5B 7A-12D+ Mc WRLr | Tern, Sooty | Sa FVvr (10, 2017) |
| Whip-poor-will, Eastern* | 3D-9D SRfc (n, ce, wc) | Plover, Snowy | FSpVacc (6, 2016) | Woodcock, American* | P SpMfc SRr FMu WRu | Bridled | Sa FVacc (2, 2008) |
| Order Apodiformes Swifts and Hummingbirds | | Wilson's | SpFVacc (3, 1998) | Snipe, Wilson's | 8A-5D+ Mc WRfc | Least* | 5A-9D+ Mu SRLu (Lc e) |
| Swift, Chimney* | 3B-10D SRc | Semipalmated | 4B-5C 7B-9D+ Mc | Sandpiper, Spotted ^o | 3D-6B 7A-12D+ Mc WVr | Gull-billed | SpVacc (1, 2009) |
| White-throated | SpFVWVacc (4, 2012) | Piping | 4B-5C 7B-10B SpMvr FMr | Solitary | 3D-5B 6B-12D+ Mfc | Caspian | 4A-6C 7A-10C+ SpMu FMfc |
| Violetear, Mexican | SFVacc (6, 2000) | Killdeer* | P c | Yellowlegs, Lesser | 2D-5D 6D-12D+ Mc WRu | Black | 4B-6B 6D-10C+ Mc |
| Hummingbird, Rivoli's | SVacc (1, 1993) | Plover, Mountain | WVacc (1, 1951) | Willet | 3D-5D 6C-9C+ Mu | Common | 4D-5D 7A-10C+ SpMr FMu |
| Ruby-throated* | 3B-11B+ SRc | Sandpiper, Upland | 3D-5C 6D-9D+ Mu | Yellowlegs, Greater | 3B-5D 7A-12D+ Mc WRu | Forster's | 7B-5C+ Mfc WRLu (s) |
| Black-chinned | 10D-3D WVvr | Whimbrel | 5B-5D+ 9A-9B SpMvr FMacc | Phalarope, Wilson's | 3C-5D 7A-10D Mfc | Royal | Sa FVvr (8, 2016) |
| Anna's | 10C-3B WVvr | Curlew, Eskimo | extirpated | Red-necked | 5B-6A 7B-11B SpMvr FMr | Skimmer, Black | Sa FVvr (8, 2008) |
| Broad-tailed | FWVacc (4, 2006) | Curlew, Long-billed | 4B-4D 10B-12B Mvr | Red | 8D-11D+ FMvr | Order Gaviformes Loons | |
| Rufous | 8B-4A FMr WVr | Godwit, Hudsonian | 4B-5D 8C-10B SpMLu FMvr | Jaeger, Pomarine | FVacc (1, 1983) | Loon, Red-throated | 10D-4A Mvr WRr |
| Calliope | FVacc (6, 2010) | Marbled | 4B-5D 7C-10A SpMvr FMr | Parasitic | 9A-12B FVvr | Pacific | 11A-4A+ Mr WRr |
| Broad-billed | FWVacc (3, 2017) | Turnstone, Ruddy | 5B-6B 7C-10B Mr | Long-tailed | FVacc (4, 2000) | Common | 10A-5D+ Mfc WRLc |
| Buff-bellied | WVacc (1, 1993) | Knot, Red | 5C-5D 8B-10A SpMacc FMvr | Murrelet, Long-billed | FVacc (1, 1980) | Yellow-billed | FWVacc (2, 2010) |
| Order Gruiformes Rails, Gallinules, and Cranes | | Ruff | 3C-5B 8A-9B SpMvr FMacc | Kittiwake, Black-legged | 11B-3A WVvr | Order Ciconiiformes Storks | |
| Rail, Yellow | 9A-5A Mvr WVacc | Sandpiper, Sharp-tailed | FVacc (1, 2016) | Gull, Sabine's | 9A-10C+ FMLr (sw, wc) | Stork, Wood | 6B-10B+ FVu (s, e) |
| Black | 9B-10B FMvr | Stilt | 3D-6A 7A-11D+ Mfc | Bonaparte's | 10A-4D+ Mfc WRfc | Order Suliformes Frigatebirds, Boobies, Cormorants, and Anhinga | |
| King* | P Mr SRLr (cc, s) | Curlew | FVacc (2, 2004) | Black-headed | 11C-1B WVvr | Frigatebird, Magnificent | 9B-10D+ Sa FVvr |

Reporting Bird Sightings

Ornithology has benefitted a great deal from the contributions of amateurs over the years. Birders who faithfully report significant observations and make them a part of the permanent record help ornithologists better understand the intricacies of bird distribution and abundance. Although it isn't always obvious whether a sighting should be reported, a good rule to follow is "If in doubt, make the report." More specific instances in which a report should be made include:

- If a species appears outside of its normal season or geographical region, it should be reported.
- If unexpectedly large or small numbers of a certain species are encountered, that count of individuals should be reported.
- If an unusual or unlisted species is seen, report it by all means, but keep in mind that the greater the rarity the more thorough supporting documentation should be.

Written and/or photographic documentation is required when reporting rarities or species observed out of season. The Arkansas Bird Records Committee reviews these reports. The basic information needed for all reports includes: common name, age (adult, immature, etc.), plumage (breeding, non-breeding, etc.), number seen, location (county, nearest town, etc.), and date. The more unusual reports require additional information such as time seen, light conditions, optical equipment, distance to bird and how measured, as well as a description of the bird's size, shape, color pattern, vocalizations if any were heard, and behavior. It is also important to eliminate any similarly appearing species and discuss prior experience with both the species being reported and any with which it might be confused. The names and addresses of additional observers should be included as well as any reference material consulted and how it influenced the description. If it is possible to obtain photographs or audio recordings, be sure to include copies with the report. **Above all else, remember that the documentation process is in place to gather additional, substantiating information and is not meant to intimidate potential contributors or question the authenticity of their reports.**

Reports of birds should be submitted online via the society's website at www.arbirds.org. If further assistance is needed, contact the Curator at arbirds@arbirds.org.

Dedication

This 2018 edition of the Arkansas Audubon Society field reference honors Dr. Kimberly Gray Smith (1948-2018). Kim came to UA-Fayetteville in 1981 and attained status as University Professor of Biological Sciences and Distinguished Professor. During his career he was a productive researcher and collaborator, with approximately 300 professional publications. Kim was deeply involved in numerous professional organizations in a variety of roles: officer, editor, meeting organizer, etc. He served as Editor in Chief of *The Auk* (2000-2004) journal of what is now American Ornithological Society. Kim presented a retirement seminar on April 5, 2018, "Life in the Fast Lane: My Life as a Community Ecologist." He ended his seminar with some advice to younger colleagues: "Be curious, be creative, challenge yourself to learn new things, learn the history of the things that interest you, take students on field trips, take students abroad," and finally, "have fun doing what you do...I did..."

The Arkansas Audubon Society was formed in May 1955 to promote the protection and wise use of our natural resources. Weekend meetings are held twice each year at locations throughout the state and are generally scheduled to coincide with the peaks of spring and fall migration. Guest speakers, workshops, and guided field trips are featured at all meetings. A quarterly newsletter is published to keep the membership informed regarding current conservation issues, noteworthy bird records, and news of members and upcoming events. Membership is open to all interested persons. For further information, please visit the society's website at www.arbirds.org

The 2018 Field Reference was compiled by Lyndal York. Document design and typesetting by Kenny Nichols. Additional editorial assistance provided by Leif Anderson, Dick Baxter, Rob Doster, Douglas A. James, Mike Modinow, Joseph C. Neal, LaDonna Nichols, Helen Parker, Herschel Raney, and Dan Scheiman.

To request 15 or more hardcopies of this field list, e-mail the Curator at arbirds@arbirds.org to make arrangements. A donation is appreciated to offset the costs of printing and mailing. To make a donation visit www.arbirds.org

| | | | |
|--|--|--|-----------------------------|
| Booby, Brown | Vacc (8, 2017) | Eagle, Golden | 9B-4B WRr |
| Gannet, Northern | SVacc (1, 2005) | Harrier, Northern* | 8B-5B+ Mfc WRfc |
| Cormorant, Neotropic ^o | 5C-12D+ Mr SRLr (sw, se) WVvr | Hawk, Sharp-shinned* | p Mfc WRu SRr (n) |
| Double-crested* | P Mc SRLu WRc | Cooper's* | P WRfc SRLu |
| Anhinga* | 3C-12C+ SRLu WVvr (s) | Goshawk, Northern | 11A-1B WVvr |
| Order Pelicaniformes Pelicans, Herons, Ibis, and Spoonbills | | Eagle, Bald* | P WRc SRu |
| Pelican, American White | P Mfc SRLu WRfc | Kite, Mississippi* | 4C-9C+ SRc (Lu wc, nw) |
| Brown | 3C-11B+ Vr (ce, s) | Hawk, Red-shouldered* | P fc |
| Bittern, American* | 3A-5D 9B-12D+ Mu WVvr | Broad-winged* | 3D-10B+ Mfc SRfc |
| Least* | 4A-10A+ Mr SRr (Lu s, e) | Swainson's* | 3C-11C+ Mu (w) SRLr (nw) |
| Heron, Great Blue* | P c | Red-tailed* | P c |
| Egret, Great* | P SRLc WRfc (ce, s) | Rough-legged | 10D-3C WVvr |
| Snowy* | 3B-11C+ Mfc SRLfc WVvr | Ferruginous | 10D-2D WVvr |
| Heron, Little Blue* | 3B-11C+ Mc SRLc WVvr | Order Strigiformes Owls | |
| Tricolored* | 4A-10C+ Mr SRLu (s) FVu (s) WVvr (sw) | Owl, Barn* | P Lu (e) Lr (w) |
| Egret, Cattle* | 3B-11B+ Mfc SRc WVr (s) | Screech-Owl, Eastern* | P fc |
| Heron, Green* | 3B-10D+ Mc SRc WVvr (s) | Owl, Great Horned* | P fc |
| Night-Heron, Black-crowned* | 3B-12B+ Mu SRLu WVvr | Snowy | 11B-1B WVvr |
| Yellow-crowned* | 3D-11A+ Mu SRLu | Burrowing | 9D-3D WVvr (e, ce) |
| Ibis, White* | 3D-11A+ SRLc (s, sw) FVLc (s, ce) WRvr (sw) | Barred* | P fc |
| Glossy ^o | 4B-5D 7C-9D+ Mr | Long-eared | 11A-5D+ WRvr |
| White-faced | 4B-6A 8A-11D+ Mr WVvr (se) | Owl, Short-eared | 10C-4C WRLu |
| Spoonbill, Roseate | 7A-10D+ FVu (ce, s) | Northern Saw-whet | 11A-2D WVr |
| Order Cathartiformes Vultures | | Order Coraciiformes Kingfishers | |
| Vulture, Black* | P fc (w) r (e) | Kingfisher, Belted* | P c |
| Turkey* | P c | Order Piciformes Woodpeckers | |
| Order Accipitriformes Kites, Eagles, and Hawks | | Woodpecker, Lewis's | SVacc (1, 1967) |
| Osprey* | P Mfc SRLr WRr (s) | Red-headed* | P fc |
| Kite, White-tailed | 3A-4D+ SpWVvr | Red-bellied* | P c |
| Swallow-tailed ^o | 4A-9D Mvr SRLvr | Sapsucker, Yellow-bellied | 9B-5B+ WRc |

| | | | |
|--|-------------------------------------|---|--------------------------------------|
| Woodpecker, Downy* | P c | Flycatcher, Least | 4C-5D 8B-10B SpMfc FMu |
| Red-cockaded* | P Lr (ec, se, wc) | Phoebe, Eastern* | P SRc SRr (se) WRfc |
| Hairy* | P fc | Say's | 9C-4D WVr |
| Flicker, Northern* | P SRu WRfc | Flycatcher, Vermilion | 10A-4B+ WVvr |
| Woodpecker, Pileated* | P fc | Family Laniidae Shrikes | |
| Ivory-billed | Formerly P Recent status unclear | Shrike, Loggerhead* | P SRu (Lfc e, nw) WRfc |
| Order Falconiformes Falcons | | Northern | WVvr (2, 2011) |
| Caracara, Crested | SpVacc (2, 2017) | Family Vireonidae Vireos | |
| Kestrel, American* | P Mc SRu WRc | Vireo, White-eyed* | 3C-11D+ SRc WVvr (s) |
| Merlin | 9B-4C+ Mu WVu | Bell's* | 4C-9D Mu SRu |
| Falcon, Peregrine° | 8C-5B+ Mu WVr | Yellow-throated* | 3D-10C+ SRfc |
| Prairie | 10C-2B+ WVr | Blue-headed | 9A-5D+ Mfc WRr (s) |
| Order Psittaciformes Parakeets | | Philadelphia | 4C-5D 9B-10B+ Mfc |
| Parakeet, Carolina | extinct | Warbling* | 4B-9D+ SRLu Mfc |
| Order Passeriformes Family Tyrannidae Flycatchers | | Red-eyed* | 4A-10B+ SRc |
| Flycatcher, Ash-throated | FWVacc (2, 2010) | Family Corvidae Jays and Crows | |
| Great-crested* | 4B-10A+ SRc | Jay, Blue* | P c |
| Kingbird, Couch's | WVacc (1, 1995) | Nutcracker, Clark's | SpVacc (1, 1891) |
| Cassin's | SpSVacc (2, 1967) | Crow, American* | P c |
| Western* | 3D-10C Mu SRLu | Fish* | P SRc WRLfc (s) |
| Eastern* | 3D-10D SRc | Family Alaudidae Larks | |
| Gray | SpVacc (1, 2007) | Lark, Horned* | P WRc SRLu |
| Flycatcher, Scissor-tailed* | 3D-11D+ SRc | Family Hirundinidae Swallows | |
| Fork-tailed | WVacc (1, 1995) | Martin, Purple* | 2A-9B+ SRc |
| Flycatcher, Olive-sided | 4D-6D 8A-10B Mu | Swallow, Tree* | 2C-11C+ Mc SRLfc |
| Wood-Pewee, Eastern* | 4A-10D SRc | Northern Rough-winged* | 3A-10D+ SRc |
| Flycatcher, Yellow-bellied | 4D-5D 8B-10C SpMr FMvr | Bank* | 4A-5D 7A-9D+ Mfc SRvr |
| Acadian* | 4A-9D+ SRfc | Cliff* | 3B-10B+ Mc SRc |
| Alder | 5A-6A 8A-9B+ Mu | Cave | 3D-5D 6D-10A+ SpVvr SFMu (sw, se) |
| Willow* | 5A-9C SpMu SRLr (nw) | Barn* | 3A-11D+ SRc |

| | | | |
|--|-----------------------------|---|---------------------------|
| Family Paridae Chickadees and Titmice | | Thrush, Swainson's | 4A-5D 9A-10C+ SpMc FMu |
| Chickadee, Carolina* | P c | Hermit | 10A-4D+ WRc |
| Titmouse, Tufted* | P c | Wood* | 4A-10B+ SRfc |
| Family Sittidae Nuthatches | | Robin, American* | P c |
| Nuthatch, Red-breasted | 9B-5B+ WRirr | Thrush, Varied | WVacc (1, 2010) |
| White-breasted* | P c (n, wc) | Family Mimidae Mockingbirds, Thrashers, and Allies | |
| Brown-headed* | P c (s, wc) | Catbird, Gray* | P SRc WRr |
| Family Certhiidae Creepers | | Thrasher, Brown* | P c |
| Creeper, Brown° | 10A-4C+ WRfc | Sage | SpFVacc (3, 2018) |
| Family Troglodytidae Wrens | | Mockingbird, Northern* | P c |
| Wren, Rock | 10B-4A+ WRr (w) | Family Sturnidae Starlings | |
| House* | P Mfc WRu SRfc (nw) | Starling, European* | P c |
| Winter | 10A-4D+ WRfc | Family Bombycillidae Waxwings | |
| Sedge* | P Mfc SRLfc (e, nw) WRLu | Waxwing, Bohemian | 1D-3C+ WVvr |
| Marsh | 9C-5C Mfc WRu | Cedar* | P SRu (nw) WRc |
| Carolina* | P c | Family Passeridae Old World Sparrows | |
| Bewick's* | P SRr (nw) WRr | Sparrow, House* | P c |
| Family Polioptilidae Gnatcatchers | | Family Motacillidae Pipits | |
| Gnatcatcher, Blue-gray* | 3C-10C+ SRc WVr | Pipit, American | 9D-4C+ Mfc WRfc |
| Family Regulidae Kinglets | | Sprague's | 10A-4D Mr WRr |
| Kinglet, Golden-crowned | 9D-4D WRc | Family Fringillidae Finches | |
| Ruby-crowned | 9D-5A+ WRc | Brambling | WVacc (1, 2016) |
| Family Muscipapidae Wheatears | | Grosbeak, Evening | 11B-5A+ Witr vr |
| Wheatear, Northern | FVacc (1, 1990) | Grosbeak, Pine | WSpSVacc (4, 1989) |
| Family Turdidae Bluebirds, Thrushes, and Allies | | Rosy-Finch, Gray-crowned | SpVacc (1, 2012) |
| Bluebird, Eastern* | P c | Finch, House* | P c |
| Mountain | FSpVacc (2, 2013) | Purple° | 10D-4B+ WRfc |
| Solitaire, Townsend's | 10C-4A WRvr (wc) | Redpoll, Common | 12A-4A WVvr (n) |
| Veery | 4C-5C+ SpMu | Crossbill, Red° | 9A-7D+ Vr |
| Thrush, Gray-cheeked | 4A-5D 9B-10C+ SpMfc FMvr | White-winged | 1A-4C WSpVvr |

| | | | |
|--|------------------------------------|---|---|
| Siskin, Pine* | 10B-5B+ Mirr WRirr | Sparrow, Song | 10A-5B+ WRc |
| Goldfinch, Lesser | SpSWVacc (6, 2017) | Lincoln's | 10A-5C+ Mfc (w, ce) WRu (w) |
| American* | P WRc SRc (nw, ce) | Swamp | 10A-5A+ WRc |
| Family Calcariidae Longspurs and Snow Bunting | | White-throated | 10A-5B+ WRc |
| Longspur, Lapland | 11B-3A+ WRfc | Harris's | 10C-5B WRu (w, ce) |
| Chestnut-collared | 12D-1D+ WVvr | White-crowned | 10A-5D WRc |
| Smith's | 11A-3C Mr WRLu (e) | Junco, Dark-eyed | 10B-4B+ WRc |
| McCown's | FVacc (2, 1996) | Family Icteridae Chat | |
| Bunting, Snow | 11C-12D+ WVvr | Chat, Yellow-breasted* | 4C-9D+ SRc |
| Family Passerellidae New World Sparrows | | Family Icteridae Blackbirds and Allies | |
| Towhee, Green-tailed | FVacc (1, 1952) | Blackbird, Yellow-headed | 4A-5C+ 8C-10C+ SpMu FMr WVvr |
| Spotted | 10C-4A+ WRr | Bobolink | 4C-5D 8C-10B+ SpMfc FMLu |
| Eastern* | P SRfc WRc | Meadowlark, Eastern* | P c |
| Sparrow, Rufous-crowned* | P Lr (wc) | Western | 10C-4D+ WRu |
| Cassin's* | SpFVacc (2, 2011) | Oriole, Orchard* | 4A-9C+ SRc |
| Bachman's* | 4A-9C+ SRLu | Bullock's | 12D-2C WVvr |
| American Tree | 10B-4A Mr WRr (n) | Baltimore* | 4A-9D+ SRLfc (ce, e, s) SRLu (nw) WVvr (s) |
| Chipping* | P SRc (n, wc) WRc (s) | Scott's | WSpVacc (3, 2013) |
| Clay-colored | 4B-5B 9A-11B+ Mu (w) Mr (e) | Blackbird, Red-winged* | P c |
| Brewer's | WVacc (1, 2011) | Cowbird, Brown-headed* | P c |
| Field* | P SRfc WRc | Blackbird, Rusty | 10C-4C+ WRfc |
| Vesper | 10A-4D Mfc WRu (s, e) | Brewer's | 11A-4B+ WRfc (e, sw) WRu (nw) |
| Lark* | 3D-9D+ SRfc SRu (nw) WVvr (s) | Grackle, Common* | P c |
| Bunting, Lark | WSpVacc (5, 2017) | Great-tailed* | P Lu (w) |
| Sparrow, Savannah | 10A-5C+ WRc | Family Parulidae Wood Warblers | |
| Grasshopper* | 4A-11D+ SRLu SRLfc (nw) | Ovenbird* | 4B-10B+ SRc (n, ce, wc) |
| Henslow's* | P Mr SRLvr (nw, n, wc) WRr (se) | Warbler, Worm-eating* | 4A-9D+ SRu |
| Le Conte's | 10A-5B WRLfc | Waterthrush, Louisiana* | 3C-10B SRfc |
| Nelson's | 5B-5C 9D-10D+ Mr | Northern | 4B-5C 8B-10A+ SpMc FMfc |
| Fox | 10B-4C+ WRfc | Warbler, Bachman's | extinct |

| | | | |
|------------------------|---|--|--------------------------------|
| Warbler, Golden-winged | 4D-5B 8D-10B Mu | Warbler, Black-throated Green* | 3D-10C+ Mc SRLu (nc, nw) |
| Blue-winged* | 4A-10B Mfc SRLu (n, ce) | Canada | 4D-6A 8D-10A Mu |
| Black-and-white* | 3C-10B+ SRc WVvr (s) | Wilson's | 4D-6A 8D-10A+ Mu (e) Mc (w) |
| Prothonotary* | 4A-9C+ SRfc | Tanagers, Cardinal, and Grosbeaks Family Cardinalidae | |
| Swainson's* | 4A-9B+ SRLu | Tanager, Summer* | 4B-10B+ SRc WVvr |
| Tennessee | 4B-5C 9A-10C+ Mc | Scarlet* | 4B-10B+ Mfc SRfc (n, wc) |
| Orange-crowned | 9D-5C+ Mfc WRu | Western | 8D-5B WVvr |
| Nashville | 4A-5C 9A-10D+ Mc | Cardinal, Northern* | P c |
| Connecticut | 4C-5D SpMvr | Grosbeak, Rose-breasted | 4C-5C 9C-10C+ SpMfc FMu |
| Mourning | 4D-5D 8D-9D+ Mu | Black-headed | 9D-5B+ WVvr |
| Kentucky* | 4B-9D+ SRfc | Blue* | 4B-10B SRc |
| Yellowthroat, Common* | 4A-10D+ SRc WRu (s, e) | Bunting, Lazuli | 2D-6B SpVvr |
| Warbler, Hooded* | 4A-10A+ SRfc | Indigo* | 4B-11A+ SRc WVvr |
| Redstart, American* | 4B-10C Mfc SRu (Lfc nw) | Painted* | 4B-9D+ SRfc |
| Warbler, Cape May | 4D-5C 8D-11D+ Mr | Dickcissel* | 4D-10B+ SRc WVvr |
| Cerulean* | 4C-9A+ SRu (Lfc nw) | | |
| Parula, Northern* | 3C-10C+ SRc | | |
| Warbler, Magnolia | 4B-5D+ 9A-10B+ Mfc Mc (e) | | |
| Bay-breasted | 4B-6B 9D-10B+ Mu | | |
| Blackburnian | 4D-6A 9A-10B Mu Mfc (e) | | |
| Yellow* | 4B-9D+ Mfc SRLu (nw) | | |
| Chestnut-sided* | 4D-10B+ SpMfc FMfc (e) u (w) SRLr (nc, nw) | | |
| Blackpoll | 4B-5D 10B-10D SpMfc FMvr | | |
| Black-throated Blue | 4C-5B 9C-10C+ Mvr | | |
| Palm | 4C-5A 9C-11C+ Mfc WRr | | |
| Pine* | P c WRu (nw) | | |
| Yellow-rumped | 9D-5B+ WRc | | |
| Yellow-throated* | 3D-10C+ SRfc | | |
| Prairie* | 4B-9C+ SRfc | | |
| Black-throated Gray | FVacc (1, 2005) | | |